

IN THE CLAIMS

This listing of the claims replaces all prior versions of the claims in the application.

1. (Withdrawn) An isolated polypeptide selected from the group consisting of:
 - a) a polypeptide comprising an amino acid sequence of SEQ ID NO:1,
 - b) a polypeptide comprising a naturally occurring amino acid sequence having at least 90% sequence identity to the amino acid sequence of SEQ ID NO:1, and
 - c) a biologically active fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:1.
2. (Withdrawn) An isolated polypeptide of claim 1 comprising an amino acid sequence of SEQ ID NO:1.
3. (Currently Amended) An isolated polynucleotide encoding a polypeptide ~~of claim 1~~ selected from the group consisting of:
 - a) a polypeptide comprising an amino acid sequence of SEQ ID NO:1,
 - b) a polypeptide comprising a naturally occurring amino acid sequence having at least 90% sequence identity to the amino acid sequence of SEQ ID NO:1, said polypeptide having phosphatase inhibitory activity, and
 - c) a biologically active fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:1, said fragment having phosphatase inhibitory activity.
4. (Original) An isolated polynucleotide encoding a polypeptide of claim 2.
5. (Previously Presented) An isolated polynucleotide of claim 4 comprising the sequence of SEQ ID NO:3.
6. (Original) A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.

7. (Original) A cell transformed with a recombinant polynucleotide of claim 6.
8. (Canceled)
9. (Original) A method for producing a polypeptide of claim 1, the method comprising:
- culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and
 - recovering the polypeptide so expressed.
10. (Withdrawn) An isolated antibody which specifically binds to a polypeptide of claim 1.
11. (Currently Amended) An isolated polynucleotide selected from the group consisting of:
- a polynucleotide comprising a polynucleotide sequence ~~selected from the group consisting~~ of SEQ ID NO:3,
 - a polynucleotide comprising a naturally occurring polynucleotide sequence having at least 90 ~~70~~% sequence identity to the polynucleotide sequence ~~selected from the group consisting~~ of SEQ ID NO:3,
 - a polynucleotide complementary to a polynucleotide of a),
 - a polynucleotide complementary to a polynucleotide of b), and
 - an RNA equivalent of a)-d).
12. (Canceled)
13. (Withdrawn) A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:
- hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample,

and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and

- b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

14. (Canceled)

15. (Canceled)

16. (Withdrawn) A composition comprising an effective amount of a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

17. (Withdrawn) A composition of claim 16, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO:1.

18. (Canceled)

19. (Withdrawn) A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting agonist activity in the sample.

20. (Canceled)

21. (Canceled)

22. (Withdrawn) A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting antagonist activity in the sample.

23. (Canceled)

24. (Canceled)

25. (Withdrawn) A method of screening for a compound that specifically binds to the polypeptide of claim 1, said method comprising the steps of:

- a) combining the polypeptide of claim 1 with at least one test compound under suitable conditions, and
- b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a compound that specifically binds to the polypeptide of claim 1.

26. (Canceled)

27. (Canceled)

28. (Withdrawn) A method for assessing toxicity of a test compound, said method comprising:

- a) treating a biological sample containing nucleic acids with the test compound;
- b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide of claim 11 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 11 or fragment thereof;
- c) quantifying the amount of hybridization complex; and
- d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

29-60. (Canceled)

61. (Previously Presented) A microarray wherein at least one element of the microarray is a polynucleotide of claim 11.

62. (Previously Presented) A method of generating an expression profile of a sample which contains polynucleotides, the method comprising:

- a) labeling the polynucleotides of the sample,
- b) contacting the elements of the microarray of claim 29 with the labeled polynucleotides of the sample under conditions suitable for the formation of a hybridization complex, and
- c) quantifying the expression of the polynucleotides in the sample.